



TECHNICAL UNIVERSITY OF MOMBASA

FACULTY OF APPLIED AND HEALTH SCIENCES

DEPARTMENT OF MATHEMATICS AND PHYSICS

UNIVERSITY EXAMINATION FOR:

AMA 5106: TEST OF HYPOTHESIS

END OF SEMESTER EXAMINATION

SERIES: MAY 2016

TIME: 3 HOURS

DATE: MAY

Instructions to Candidates

You should have the following for this examination

-Answer Booklet, examination pass and student ID

This paper consists of five questions. Attempt QUESTION ONE and any other TWO.

Do not write on the question paper.

Question ONE

- State and prove Neyman-Pearson Lemma (8 marks)
- A manufacturer is interested in the output voltage of a power supply used in a PC. Output voltage is assumed to be normally distributed, with standard deviation 0.25 Volts, and the manufacturer wishes to test $H_0; \mu = 5$ Volts against $H_1; \mu \neq 5$ Volts, using 8 units.
 - The acceptance region is $4.85 \leq \bar{x} \leq 5.15$ Find the level of significance. (4 marks)
 - Find the power of the test for detecting a true mean output voltage of 5.1 Volts. (5 marks)
- Show that the class of all test functions is a convex function (3 marks)
- Define the power function of a test (4 marks)
- Show that 1-parameter exponential family has a monotone likelihood ratio. (6 marks)

Question TWO

- Let x be a random variable with probability density function $f(x)$. Find a size Γ test for; (7 marks)

$$H_0; f(x) = f_0(x) = \frac{1}{\sqrt{2f}} e^{-\frac{x^2}{2}}$$

$$H_1; f(x) = f_1(x) = \frac{1}{f} \frac{1}{1+x^2}$$

- b. Let x_1, x_2, \dots, x_n be independently identically distributed $N(0, \tau^2)$ random variables. Determine whether there exists a uniform most powerful test for the hypothesis of the form $H_0; \tau^2 = \tau_0^2$ against $H_1; \tau^2 = \tau_1^2$ (8 marks)
- c. Show that for testing $H_0; \mu_1 \leq \mu \leq \mu_2$ against $H_1; \mu < \mu_1$ or $\mu > \mu_2$ there exists a uniform most powerful unbiased size Γ test given by $W(x) = \begin{cases} 1 & \text{if } T(x) > c_1 \\ \epsilon & \text{if } T(x) = c_2 \\ 0 & \text{if } c_1 < T(x) < c_2 \end{cases}$ (5 marks)

Question THREE

- a. Define an unbiased test (5 marks)
- b. If the *pdf* $f(x; \mu)$ are such that the power function of every test is continuous and if W_0 is uniform most powerful among all tests satisfying some conditions and is level Γ test, then show that W_0 is unbiased. (5 marks)
- c. Let $X \sim \text{bin}(n, p)$, find an unbiased size Γ test for $H_0; p = p_0$ against $H_1; p = p_1$ (10 marks)

Question FOUR

- a. Let x_1, x_2, \dots, x_n be independently identically distributed $N(\mu, \tau^2)$ random variables, Let y_1, y_2, \dots, y_n be independently identically distributed $N(\mu, \tau^2)$ random variables. Where τ^2 is common. Suppose X_i 's and Y_i 's are independent. Determine a size Γ LRT test for $H_0; \mu = \mu_0$ against $H_1; \mu \neq \mu_0$ (10 marks)
- b. Let $x_{i1}, x_{i2}, \dots, x_{in}$ be independent normally distributed random variables with mean μ_i and variance τ_i^2 . Determine a Γ likelihood ratio test for the hypothesis of the form $H_0; \tau_i^2 = \tau_j^2$ against $H_1; \tau_i^2 \neq \tau_j^2$ (10 marks)

Question FIVE

- a. Determine a Γ likelihood ratio test for the hypothesis of the form $H_0; \tau^2 = \tau_0^2$ against $H_1; \tau^2 = \tau_1^2$ (μ is unknown) (10marks)
- b. Let y_1, y_2, \dots, y_n be independently identically distributed $N(S, \sigma^2)$ random variables. Find a size Γ likelihood ratio test for testing $H_0; S = S_0$ against $H_1; S \neq S_0$ (10 marks)