

TECHNICAL UNIVERSITY OF MOMBASA

FACULTY OF ENGINEERING AND TECHNOLOGY

DEPARTMENT OF ELECTRICAL & ELECTRONICS

ENGINEERING

UNIVERSITY EXAMINATION FOR:

INSTITUTION BASED

THE DEGREE IN BACHELOR OF TECHNOLOGY IN

ELECTRICAL ENGINEERING

TMC 4252 : Fluid Mechanics

END OF SEMESTER EXAMINATION

SERIES: APRIL 2017

TIME: 2 HOURS

DATE: Pick Date Select Month Pick Year

Instructions to Candidates

You should have the following for this examination -Answer Booklet, examination pass and student ID This paper consists of Choose No questions. Attempt Choose instruction. **Do not write on the question paper.**

Q.1

- a. Describe the following types of fluids:
 - (i) Ideal fluid

		(ii)	Real fluids	
		(iii)	Newtonian fluids	
		(iv)	Non-Newtonian fluids	(6 marks)
	b. Explain the following terms			
		i.	Density	
		ii.	Specific Volume	
		iii.	Specific gravity	
		iv.	Adhesion	
		٧.	Surface Tension	(5marks)
	c. A reservoir of glycerin has a mass of 1200kg and volume of 0.952m ² . find the following			I the following:
		i.	The glycerin's weight	
		ii.	The glycerin's density	
		iii.	The glycerin's specific weight	
		iv.	The glycerin's specific gravity	(6 marks)
d		The volume of a rock is found to be .00015m ³ . If the rock's specific gravity is 2.60. What is		
		weight?		(3marks)
Q.2				
	a.	Derive the equation for the relationship between gauge pressure P inside a droplet of		
		liquid and surface tension. (4marks)		
	b.	The pressure outside the droplet of water of diameter 0.04mm is 10.32		

e aropiet o ater of diameter 0.04r n is 10.3 N/cm²(atmospheric pressure).Calculate the pressure within the droplet if the surface (4 marks) tension is given as 0.0725N/m of water.


c. Explain the phenomenon of capillary (2 marks)

d. Find the minimum size of glass tube that can be used to measure water level if the capillary rise in the tube is to be restricted to 2mm. Consider the surface tension of water in contact with air as 0.07375N/m (4 marks)

e. A U-tube is made up of two capillaries of bores 1.2 m and 2.4 mm respectively. The tube is held vertical and partially filled with liquid of surface tension 0.06 N/m and zero contactangle. If the estimated difference in the level of two menisci is 15 mm, determine the mass density of the liquid. (6 marks)

Q.3

- a. Benzene flows through a 100mm diameter pipe at a mean velocity of 3.00m/s. Find the following
 - i. Volume flow rate in m³/s
 - ii. Mass flow rate Use (density of Benzene 879kg/m³) (4 marks)
- b. A pipe (1) 450 mm in diameter branches into two pipes (2 and 3) of diameters300 mm and 200 mm respectively as shown in Fig. 1. If the average velocity in 450 mm diameter pipe is 3 m/s find:
 - i. Discharge through 450 mm diameter pipe;
 - ii. Velocity in 200 mm diameter pipe if the average velocity in 300 mm pipe is 2.5 m/s.



- c. List the assumptions which are made while deriving Bernoulli's equation (5 marks)
- A pipe 200 m long slopes down at 1 in 100 and tapers from 600 mm diameter at the higher end to 300 mm diameter at the lower end, and carries 100 litres/sec of oil (sp. gravity 0.8). If the pressure gauge at the higher end reads 60 kN/m², Calculate:
 - i. Velocities at the two ends;
 - ii. Pressure at the lower end.

Neglect all losses. (6 marks)

(5 marks)

Q4

a. Define the following terms

- i. Total Pressure
- ii. Centre of pressure (4 marks)
- b. A rectangular plane surface is 2 m wide and 3 m deep It lies in vertical plane in water. Determine the total pressure and the centre of pressure on the plane surface when its upper edge is horizontal and coincides with the water surface. (6 marks)

Use
$$I_G = \frac{bd}{12}$$

c. Define the following terms

i. Buoyancy

- ii. Centre of buoyancy
- Find the volume of the water displaced and the position of centre of buoyancy for a wooden block of width 2.5 and of depth 1.5, when it floats horizontally in water. The wooden block 650kg/m3 and its length 6 m.

Question Five

- a) Define the following co-efficients
 - i. Co-efficient of velocity
 - ii. Co-efficient of contraction
 - iii. Co-efficient of discharge
- b) Water discharged at the rate of 98.2 litres /seconds through a 120 mm diameter vertical sharpedged orifice placed under a constant head of 10 metres. A point on the jet, measured from the vena-contracta of the jet has co-ordinates 4.5 metres horizontal and 0.54 metres vertical. Find
 - a. Co-effientient of velocity
 - b. Co-efficient of contraction
 - c. Co-efficient of discharge (6 marks)
- c) Describe the following devices used for measuring the rate of flow
 - i. Notch
 - ii. Weir
 - iii. Orifice (3marks)
- d) A rectangular notch has a discharge of 0.24 m3/s, when head of water is 800 mm. Find the length of the notch. Assume Cd = 0.6. (5 marks)

(4 marks)

(6 marks)