

Technical University of Mombasa

Faculty of Applied and Health Sciences

DEPARTMENT OF ENVIRONMENT AND HEALTH SCIENCES

CERTIFICATE IN COMMUNITY HEALTH AND HIV MANAGEMENT (CCH 13)

AML 1106 MEDICAL PARASITOLOGY

SEMESTER: EXAMINATIONS
SERIES: DECEMBER 2013
TIME: 2 HOURS

INSTRUCTIONS:

You should have the following for this paper
- Answer booklet
This paper consists of section A and B.
Answer ALL question
This paper consists of 2 PRINTED pages

- 1. Toxoplasma gondi is a parasite of
 - a) Goats
 - b) Cats
 - c) Man
 - d) Cattle
- 2. Malabsoption of fats is a symptom associated with
 - a) Giardiasis
 - b) Nacaroriasis
 - c) Teaniasis
 - d) Turgosis
- 3. Conjugation is a mode of reproduction with
 - a) Balantidium coli
 - b) Entamoeba gingivalis
 - c) Trichomonus vaginalis
 - d) Trypanosonus
- 4. "Free falling leaf" movement is used to describe the movement of
 - a) Ascaris lumbricoides
 - b) Giadia lamblia
 - c) Trichomonus vaginalis
 - d) Trichomonus gingivalis
- 5. A large circular karyosome sorounded by refractive globules belong to
 - a) Iodamoeba butschilli
 - b) Iodamoeba fragilis
 - c) Entamoeba coli
 - d) Entamoeba histolytica
- 6. Flukes associated with sheep rearing is
 - a) Heterophyles heterophyles
 - b) Schistosomes haematobium
 - c) Schistosomes haematobium
 - d) Paragonimus westermani
- 7. Which coastode does not require an intermediate host in its life cycle
 - a) Hymenolepis nana
 - b) Hymenolepis diminura
 - c) Taenia solium
 - d) Echinococus granulosus

- 8. The genus Trichomonus have the feature
 - a) Undularting membrane
 - b) Psedopodia
 - c) Cilite
 - d) Trophozoite
- 9. The largest human trematode is
 - a) Heterophyes heterophyes
 - b) Fasciolopsis buski
 - c) Fasciola hepatica
 - d) Schistosomes
- 10. Lancet fluke is
 - a) Dicrocoelium dentriticum
 - b) Heterophyes hererophye
 - c) PAragonimus westermani
 - d) Schistosomes
- 11. Wearing of shoes is important in controlling
 - a) Netatoriasis
 - b) Schistosomiasis
 - c) Elephantiasis
 - d) Ascariasis
- 12. Transmission of the following parasite is by ingesting edible acquatic plants
 - a) Heterophyes heterophyles
 - b) Fasciola hepatica
 - c) Fasciola buski
 - d) Schistosomes
- 13. Cysts with splineter shaped chromatoidal bars belong to
 - a) Entamoeba histolytica
 - b) Entamoeba coli
 - c) Trichomonus vaginalis
 - d) Trichomonus
- 14. Coadrin-china diarrhea is synonymous to
 - a) Ascariasis
 - b) Strongyloidiasis
 - c) Nacatonasis
 - d) Whip worm infection

- 15. Infection with schistosome is through
 - a) Ingestion of eggs
 - b) Skin penetration
 - c) Innoculation
 - d) Blood transfusion
- 16. Possession of genital sucker is a characteristic of
 - a) Heterophyes heterophyes
 - b) Schistosomes
 - c) Paragonimus westermani
 - d) Opistmorcins sinensis
- 17. The egg of the following flukes have a leveral spine
 - a) Schistosoma japonicum
 - b) Schistosoma manson
 - c) Heterophyes heterophyes
 - d) Paragonimus westemani
- 18. The Chinese liver fluke is synonymous to
 - a) Dicrocoehum dentriticum
 - b) Chronorchis sinensis
 - c) Opisthochis viverini
 - d) Schostosomes
- 19. The following is not a characteristic of amoeba.
 - a) Move by means of pseudopodia
 - b) Reproduction is by simple binary fussion
 - c) They form cysts
 - d) Have cytoplasm divided into granular ectoplasm and clear endoplasm.
- 20. The following are organelles of locomotion protozoa
 - a) Cillia and Karosome
 - b) Flagella and cilia
 - c) Pseudopodia, vacudes and flagella
 - d) Pseudopodia and chromatin granules
- 21. The following protozoa trophozoite has 2 nucleus but lacks median body
 - a) Giadia lamblia
 - b) Entamoebacoli
 - c) Trichomonus vaginalis
 - d) Dientamoeba fragilis

- 22. Chagas disease is also called
 - a) South American disease
 - b) Trypanosomiasis
 - c) South American Trypanesomicidis
 - d) South African trypanosomiasis
- 23. The following are of tissue flagellases except
 - a) Crithidia
 - b) Leishmania form
 - c) Leptomonad
 - d) Mastigote
- 24. The following are modes of infection of trypanosome bruzi cruzi
 - a) Contaminative
 - b) Congenital
 - c) Orality
 - d) Innoculation
- 25. New dehhi boil is associated with
 - a) Leishmania donovani
 - b) Leishmania tropica
 - c) Leishmmania braciliensis
 - d) LEishanania guynensi
- 26. Habitat of Babesia is
 - a) RBC
 - b) WBC
 - c) Platelates
 - d) Plasma
- 27. The following parasites are tranmited by inoculation except
 - a) Plasmodium
 - b) Leishmania
 - c) Babesia
 - d) Isospora belli
- 28. The following reagents are used in modified ritches concentration except
 - a) Formaline
 - b) Saline
 - c) Ether
 - d) Eosine

- 29. The following are non pathogenic flagellates except
 - a) Enteromonas Homin's
 - b) Trichomonas hominis
 - c) TRichomonas tenax
 - d) Trichomanos vaginalis
- 30. Inhalation is a mode of infection with
 - a) Strongyloides stercoralis
 - b) Enterobius vermicularis eggs
 - c) Hookworms
 - d) Ascaris lumbricoides adult worm
- 31. The eggs of the following parasite are flattened on the ventral side and convex on the dorsal side
 - a) Strongyroides stercoralis
 - b) Hookworms
 - c) Trichuris trichiura
 - d) Enterobius vermicularis
- 32. Heterogonic and hologonic phases are associated with the following parasites
 - a) Wuchereria bancrofti
 - b) Brugia malayi
 - c) LoaLoa
 - d) Strongyloides stercoralis
- 33. The following larvae have invaginated stolex
 - a) Procecoide larvae
 - b) Plerocercoide larvae
 - c) Echinococcus
 - d) Filariform larvae
- 34. Cysticercus boris is
 - a) The larvae of Teania serginata
 - b) The larvae of Taenia solium
 - c) The larvae of echinococus granulosus
 - d) The larvae of hemimolepis nana
- 35. The longest tapeworm infecting man is
 - a) Dypilidium caninum
 - b) Diphylobothrium latum
 - c) Taenia solium
 - d) Taenia serginata

- 36. The following is a genera of pseudophyllidian cestode
 - a) Spirometra
 - b) Eshinococcus
 - c) Dypilidium
 - d) Hymenolepis
- 37. Sparganosis is associated with
 - a) Hymenlepis deminuta
 - b) Taenia serginata
 - c) Taena solium
 - d) Spirometra mansoni
- 38. The following flukes do not encyst
 - a) Schistosomes
 - b) Fasciola
 - c) Fasciolopsis
 - d) Dicrocoelum
- 39. Body boses is a features of the parasite
 - a) LoaLoa
 - b) Onchocerca
 - c) Wuchereria
 - d) Mansonella
- 40. Guinea worm is synonymous to
 - a) Dracunculus medinensis
 - b) Mansonella ozzandi
 - c) Wucherenia bancrofi
 - d) Ascaris lumbricoides

SECTION B Essay

1. Discuss the effects of parasites to the host (20marks)

2. Using well labelled diagram describe the different types of cercaria of trematodes. (20marks)

- 3. (i) Discuss the control measures of the following
 - (a) Teania saginates

(5marks)

(b) Schistosomes

(5marks)

- (ii) Differentiate the following and give examples
 - (a) Intermediate and deffitive host
 - (b) Endoparasite and ectoparasite
 - (c) Obligate and facultative parasite
 - (d) Vector and parasite

(10marks)